

The Mean Value Theorem

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the value or values of c that satisfy the equation $\frac{f(b) - f(a)}{b - a} = f'(c)$ in the conclusion of the Mean Value Theorem

for the function and interval.

1) $f(x) = x^2 + 3x + 1$, $[-1, 2]$

A) $-1, 2$

B) $\frac{1}{2}$

C) $0, \frac{1}{2}$

D) $-\frac{1}{2}, \frac{1}{2}$

1) _____

2) $f(x) = x + \frac{12}{x}$, $[3, 4]$

A) $0, 2\sqrt{3}$

B) $2\sqrt{3}$

C) $3, 4$

D) $-2\sqrt{3}, 2\sqrt{3}$

2) _____

3) $f(x) = \ln(x - 1)$, $[2, 4]$

A) 2.820

Round to the nearest thousandth.

B) 3.820

C) 3.885

D) ± 2.820

3) _____

4) $f(x) = x^2 + 3x + 4$, $[1, 2]$

A) 1, 2

B) $-\frac{3}{2}, \frac{3}{2}$

C) $\frac{3}{2}$

D) $0, \frac{3}{2}$

4) _____

5) $f(x) = x + \frac{18}{x}$, $[2, 9]$

A) $-3\sqrt{2}, 3\sqrt{2}$

B) 2, 9

C) $0, 3\sqrt{2}$

D) $3\sqrt{2}$

5) _____

6) $f(x) = \ln(x - 1)$, $[2, 5]$

A) 4.164

Round to the nearest thousandth.

B) 3.164

C) ± 3.164

D) 3.731

6) _____

7) $f(x) = x^2 + 2x + 4$, $[1, 2]$

A) 1, 2

B) $\frac{3}{2}$

C) $0, \frac{3}{2}$

D) $-\frac{3}{2}, \frac{3}{2}$

7) _____

8) $f(x) = x^2 + 5x + 4$, $[-3, 2]$

A) $0, -\frac{1}{2}$

B) $-\frac{1}{2}, \frac{1}{2}$

C) $-\frac{1}{2}$

D) $-3, 2$

8) _____

9) $f(x) = \ln(x - 4)$, $[5, 9]$

A) 6.885

Round to the nearest thousandth.

B) 6.485

C) 7.485


D) ± 6.485

9) _____

Find the open intervals on which the function is increasing and decreasing. Identify the function's local and absolute extreme values, if any, saying where they occur.

10)


10) _____



- A) increasing on $(-3, 3)$; decreasing on $(-6, 0)$;
absolute maximum at $(3, 3)$; absolute minimum at $(-3, -3)$
- B) increasing on $(-3, 3)$; decreasing on $(-6, -3)$ and $(3, 6)$;
absolute maximum at $(3, 3)$; absolute minimum at $(-3, -3)$
- C) increasing on $(-3, 3)$; decreasing on $(0, 6)$;
absolute maximum at $(3, 3)$; absolute minimum at $(-3, -3)$
- D) increasing on $(-3, 3)$; decreasing on $(-6, -3)$ and $(3, 6)$;
no absolute maximum; no absolute minimum

11)

11) _____



- A) increasing on $(-2, 0)$ and $(2, 4)$; decreasing on $(0, 2)$;
absolute maximum at $(4, 6)$; local maximum at $(0, 2)$; absolute minimum at $(-2, 0)$ and $(2, 0)$
- B) increasing on $(-2, 0)$ and $(2, 4)$; decreasing on $(0, 2)$;
absolute maximum at $(4, 6)$; absolute minimum at $(-2, 0)$ and $(2, 0)$
- C) increasing on $(2, 4)$; decreasing on $(0, 2)$;
absolute maximum at $(4, 6)$; local maximum at $(0, 2)$; absolute minimum at $(-2, 0)$ and $(2, 0)$
- D) increasing on $(-2, 0)$ and $(2, 4)$; decreasing on $(0, 2)$;
absolute maximum at $(4, 6)$ and $(0, 2)$; absolute minimum at $(-2, 0)$ and $(2, 0)$

Answer Key

Testname: THE MEAN VALUE THEOREM

- 1) B
- 2) B
- 3) A
- 4) C
- 5) D
- 6) B
- 7) B
- 8) C
- 9) B
- 10) B
- 11) A